

Geophysical Research Letters®

RESEARCH LETTER

10.1029/2023GL102815

Key Points:

- Terrestrial carbon sinks estimated based on GOSAT XCO₂ and net ecosystem exchanges (NEE) from 12 terrestrial biosphere models using atmospheric inverse method
- The uncertainty reduction rates decrease significantly at decreasing spatial scales
- The GOSAT XCO₂ can only promise a robust continental-scale NEE estimate

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

F. Jiang,
jjiangf@nju.edu.cn

Citation:

Zhang, L., Jiang, F., He, W., Wu, M., Wang, J., Ju, W., et al. (2023). A robust estimate of continental-scale terrestrial carbon sinks using GOSAT XCO₂ retrievals. *Geophysical Research Letters*, 50, e2023GL102815. <https://doi.org/10.1029/2023GL102815>

Received 10 JAN 2023
 Accepted 11 MAR 2023

© 2023. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](https://creativecommons.org/licenses/by/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

A Robust Estimate of Continental-Scale Terrestrial Carbon Sinks Using GOSAT XCO₂ Retrievals

Lingyu Zhang¹, Fei Jiang^{1,2,3} , Wei He¹ , Mousong Wu¹ , Jun Wang¹ , Weimin Ju^{1,2,3} , Hengmao Wang¹ , Yongguang Zhang^{1,2,3} , Stephen Sitch⁴ , Anthony P. Walker⁵ , Xu Yue⁶ , Shuzhuang Feng¹, Mengwei Jia¹, and Jing M. Chen⁷ 

¹Jiangsu Provincial Key Laboratory of Geographic Information Science and Technology, International Institute for Earth System Science, Nanjing University, Nanjing, China, ²Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing, China, ³Frontiers Science Center for Critical Earth Material Cycling, Nanjing University, Nanjing, China, ⁴College of Life and Environmental Sciences, University of Exeter, Exeter, UK, ⁵Environmental Sciences Division and Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN, USA, ⁶School of Environmental Science and Engineering, Nanjing University of Information Science & Technology (NUIST), Nanjing, China, ⁷Department of Geography and Planning, University of Toronto, Toronto, Ontario, Canada

Abstract Satellite XCO₂ retrievals could improve the estimates of surface carbon fluxes, but it remains unknown on what scales these estimates are robust. Here, we use the time-dependent Bayesian synthesis top-down method and prior net ecosystem exchanges (NEEs) from 12 terrestrial biosphere models (TBMs) to infer the monthly carbon fluxes of 51 land regions with constraints by GOSAT XCO₂ retrievals. We find that the uncertainty (standard deviation of 12 TBMs) reduction rates (uncertainty reduction rate (URR)) decrease significantly at decreasing spatial scales. On the continental-scale, the mean URR is about 57%, and the annual and seasonal cycle estimates of NEE are rather robust. The evaluation shows that the posterior CO₂ concentrations are significantly improved at the continental scale. Our study suggests that the GOSAT XCO₂ can only promise a robust continental-scale NEE estimate, and improving the XCO₂ accuracy is an effective way to achieve robust estimates on smaller scales under current spatial coverage.

Plain Language Summary Remotesensing-based CO₂ measurement can improve the estimates of surface carbon fluxes due to its relatively well global coverage, but it remains unknown on what spatial scales the satellite observation could provide a robust estimate. Here, net ecosystem exchanges (NEEs) from 12 terrestrial biosphere models (TBMs) of 51 land regions for the period of 2011–2014 are constrained using GOSAT XCO₂ retrievals, and the uncertainty (standard deviation of 12 TBMs) reduction rates (uncertainty reduction rate (URR)) at different spatial scales are analyzed. We find that (a) from the whole globe to the mean of 51 regions, the URR decreases from 95% to 16%. (b) On the continental-scale, the mean URR is about 57%, and the annual NEEs in Asia, N. America, Europe, S. America, Africa, and Australia are estimated to be -2.07 ± 0.25 , -0.85 ± 0.09 , -0.76 ± 0.21 , -0.36 ± 0.17 , -0.36 ± 0.30 , and -0.15 ± 0.12 PgC yr⁻¹, respectively. Our study suggests that the GOSAT XCO₂ can only promise a robust continental-scale NEE estimate, and improving the XCO₂ accuracy is an effective way to achieve robust estimates on smaller scales under current satellite observing capacity.

1. Introduction

Terrestrial ecosystems and oceans absorb about half of anthropogenic carbon emissions (Friedlingstein et al., 2020), slowing down the increase of CO₂ in the atmosphere and thus mitigating climate change. Accurate estimation of terrestrial carbon sinks and sources is an indispensable step to understand the status and the potential of their roles in regulating climate change. As a major way of constraining terrestrial carbon flux estimates with observations over large scales, top-down atmospheric inversion infers carbon fluxes from atmospheric CO₂ mole fraction observations and a priori flux, which can effectively reduce the uncertainty of carbon flux estimates (Thompson et al., 2016). At the global or hemisphere scale, the carbon flux estimates from various atmospheric CO₂ inversions are in relatively good agreement, but at continental or regional scales, the agreement is greatly weakened due to errors in either inversion methods or observational data (Baker et al., 2006; Deng & Chen, 2011).

In situ CO₂ observations have been widely used in past atmospheric CO₂ inversions (Baker et al., 2006; Deng & Chen, 2011; Gurney et al., 2002; Jiang et al., 2013; Monteil et al., 2020; Peylin et al., 2013). Due to the uneven

distribution of global surface CO₂ observations, relatively consistent results can be obtained in places where observations are densely distributed, for example, Europe and North America (N. America). However, inversion results have high uncertainty in areas with low observation density (Maksyutov et al., 2013). The uneven distribution of observations leads to great differences in the capability of inversions to constrain the land carbon cycle in different regions (Gurney et al., 2002). Remotesensing-based CO₂ measurements provide global coverage with high spatial resolutions (Baker et al., 2010). Many studies have estimated regional carbon sources and sinks using column averaged dry air mole fractions of CO₂ (XCO₂) from GOSAT and OCO-2 satellites (Baker et al., 2010; Basu et al., 2013; Byrne et al., 2020; Chevallier et al., 2014, 2019; Crowell et al., 2019; Deng et al., 2014; Houweling et al., 2015; Jiang et al., 2021; Wang et al., 2018, 2019, 2022), boosting the possibility of better constraining the carbon cycle at finer spatial scale (Byrne et al., 2019). The GOSAT intercomparison study from Houweling et al. (2015) found that robust estimates could be obtained at large global scale, but at the TRANSCOM regions, the different inversions differed significantly. Chevallier et al. (2019) compared surface-based and satellite-based inversions and found large differences in annual fluxes between GOSAT-based and surface-based estimates at subcontinental scales. In addition, although remotesensing observations provide improved spatial coverage, the choice of different XCO₂ retrieval algorithms can lead to different results (Takagi et al., 2014). Byrne et al. (2019) explored the spatial scales of interannual variability of net ecosystem exchanges (NEE) constrained using GOSAT XCO₂, giving correlations between interannual variability at different scales and multiple “proxies,” but the spatial scales at which the inversion results are robust remain unclear.

Here, we assimilate the latest GOSAT XCO₂ observations using the time-dependent Bayesian synthesis method (“Method”) to optimize terrestrial ecosystem carbon exchange (NEE) of 51 land regions (Figure S1a in Supporting Information S1) from multiple TBMs. The inversion is from May 2009 to 2014, the first 20 month-period is treated as the spin-up stage, and the inversion results from 2011 to 2014 were analyzed in this study. NEEs simulated from 12 TBMs were used as prior fluxes within the same atmospheric inversion framework and constrained with the same observations to explore on what scales the GOSAT XCO₂ retrievals can provide robust NEE estimates.

2. Methods

2.1. Inversion Method

We use the time-dependent Bayesian synthesis method (Rayner et al., 1999), and the GOSAT XCO₂ retrievals, to estimate global surface CO₂ net fluxes. The key to this method is to minimize the following cost function (Rayner et al., 1999):

$$J = \frac{1}{2}(Ms - c)^T R^{-1}(Ms - c) + \frac{1}{2}(s - s_p)^T Q^{-1}(s - s_p) \quad (1)$$

where M is the transport operator; c is the observations; s is the vector of carbon flux combined with initial well-mixed atmospheric CO₂ concentration; s_p is a priori estimation of s ; and R and Q are the uncertainties of c and s_p , respectively. By minimizing this cost function, the posterior fluxes s_{post} and their uncertainties Q_{post} could be obtained as:

$$s_{\text{post}} = (M^T R^{-1} M + Q^{-1})^{-1} (M^T R^{-1} c + Q^{-1} s_p) \quad (2)$$

$$Q_{\text{post}} = (Q^{-1} + M^T R^{-1} M)^{-1} \quad (3)$$

The global surface is separated into 69 regions, including 51 regions for land, and 18 regions for ocean (Figure S1a in Supporting Information S1). The partition scheme of land was adopted from Wang et al. (2021). The bias-corrected GOSAT ACOS V9 XCO₂ for the years 2009–2014 is adopted as observations (O'Dell et al., 2018; Taylor et al., 2022; Wunch et al., 2017), and has been re-gridded to 1° × 1° by Jiang et al. (2021) with the best quality approach (Wang et al., 2019).

Four types of a priori carbon fluxes were used in the inversion, namely terrestrial ecosystem carbon flux (NEE), ocean (OCEAN) carbon exchange, fossil fuel and cement production (FFC) carbon emissions, and biomass burning (FIRE) carbon emissions. The OCEAN flux, FFC and FIRE emissions were adopted from the product of NOAA's CarbonTracker, version 2017 (CT2017). In many offshore areas, the OCEAN fluxes are missing,

we filled them with the fluxes of 2009 simulated by the global ocean circulation and biogeochemistry model (OPA-PISCES-T) (Buitenhuis et al., 2006; Jiang et al., 2013). The prior NEE fluxes were obtained from 12 TBMs, including BEPS (Chen et al., 1999; Ju et al., 2006), CASA (Potter et al., 1993), and 10 models from TRENDYv9 (Friedlingstein et al., 2020) (i.e., CABLE-POP (Haverd et al., 2018), DLEM (Tian et al., 2015), ISAM (Meiyappan et al., 2015), LPX-Bern (Lienert & Joos, 2018), OCN (Zaehle & Friend, 2010), ORCHIDEE (Lurton et al., 2020), ORCHIDEEv3 (Vuichard et al., 2019), SDGVM (Walker et al., 2017), VISIT (Kato et al., 2013), YIBs (Yue & Unger, 2015)). BEPS is a remotesensing-based TBM, which was driven by the LAI and clumping index products from MODIS. In this study, the BEPS simulations were adopted from Jiang et al. (2021). The CASA simulations were also derived from CT2017. There are 20 TBMs in TRENDYv9 S3 simulations, we selected the simulations with spatial resolution greater than $1^\circ \times 1^\circ$.

The transport operator M is simulated using the Model for Ozone And Related chemical Tracers, version 4 (MOZART-4) (Emmons et al., 2010). The MOZART-4 model was run at a spatial resolution of approximately $2.8^\circ \times 2.8^\circ$ (128×64 grids), and 28 vertical layers. It was driven by the ERA-Interim reanalysis data obtained from the European Centre for Medium-Range Weather Forecasts (Dee et al., 2011). Using MOZART-4, we calculated the contributions of each month and each region to the XCO_2 at each grid and time. Following Jiang et al. (2013), for each month and each region, the model is continuously run for 3 years, with 1 Pg carbon emitted in the first month and no emission in the months thereafter. The spatial distribution of emissions within each land region was assigned according to the multi-year averaged net primary production (NPP), for the ocean region, no distribution was considered. The background CO_2 concentration was set to 390 ppm, which is the average concentration of April and May 2009 observed at the global background station of Mauna Loa (Ed Dlugokencky and Pieter Tans, NOAA/GML (gml.noaa.gov/ccgg/trends/)). The OCEAN flux, FFC and FIRE emissions were assumed to be prescribed, and the CO_2 concentrations contributed from these three types of fluxes were also simulated using the MOZART-4 model. Details about the calculation of the XCO_2 contributions from each month and each region is given in Text S1 in Supporting Information S1. In order to save computational costs and reduce the size of the transport matrix M , the observations and the variables corresponding to the observations were rescaled to a resolution of $15^\circ \times 15^\circ$ per month in this study.

For the uncertainties of prior fluxes, we assumed a global land uncertainty of 2.0 PgC yr^{-1} , which was distributed to different regions based on a multi-year average annual NPP from the CASA model (Potter et al., 1993). Considering that NPP is very small in winter and large in summer, assigning uncertainty exactly according to the monthly variation in NPP would result in little uncertainty in winter, so we adopted the scheme of averaging NPP with and without monthly variation and using this result to assign uncertainty. In addition, we fixed the lowest monthly uncertainty of each region to 0.1 PgC . The annual uncertainty of global land is within the range of previous studies (Baker et al., 2006; Basu et al., 2013; Deng & Chen, 2011; Houweling et al., 2004; Rodenbeck et al., 2003). We neglected the temporal and spatial correlation of the prior flux uncertainties. According to Wang et al. (2019), the observation error over a $15^\circ \times 15^\circ$ grid for each month is 1.9 times of the average of the retrieval errors provided by the GOSAT product within that grid, but its minimum value was set to 1 ppm (Detailed in Text S2 in Supporting Information S1). For the inversion results, May 2009–December 2010 is taken as the spin-up phase, and only the inversion results from 2011 to 2014 are analyzed and discussed.

We performed four sensitivity experiments using different a priori flux uncertainty (Case Q), observation error settings (Case R), and observations aggregations (Case A10 and Case A5). For Case Q, the prior uncertainty of each land region was set to be the standard deviation of the 12 prior NEEs (Philip et al., 2019). Case R ignores the difference in observation errors, setting the observation error uniformly at 0.5 ppm, which may be the accuracy goal for future satellite observations (Sierk et al., 2021), and then scaling them up by a factor of 1.9. To investigate the impact of observations number on the inversion results, Case A10 and Case A5 were conducted, in which the XCO_2 observations were aggregated into $10^\circ \times 10^\circ$ and $5^\circ \times 5^\circ$ grids, respectively (Detailed in Table S1 in Supporting Information S1).

2.2. Evaluation Data and Method

In this study, surface CO_2 observations from the CO_2 GLOBALVIEWplus v7.0 ObsPack data set (Cox et al., 2021) were used for independent evaluations. We selected 168 sets of discrete (flask), and quasi-continuous (in situ) measurements at surface and tower with observations start date earlier than 2011, and stop date later than 2014. Of these, there are 34, 37, 75, 4, 9, and 9 sets of records available for Asia, Europe, North America, S. America,

Africa, and Oceania, respectively. In addition, in Asia, the ObsPack observations are mainly distributed in the middle and high latitudes. Therefore, we further chose the observations from the Comprehensive Observation Network for Trace gases by Airliner (CONTRAIL) project (Machida et al., 2008; Machida et al., 2019 (Reference date: 2021/10/29); Matsueda et al., 2008; Matsueda et al., 2015), to evaluate the posterior CO₂ over South-east Asia. The CONTRAIL project measures CO₂ concentrations on several Boeing aircrafts along their flight paths. Vertical profiles of CO₂ concentrations near airports were observed during the taking off and landing. We selected observations between 2,000 and 6,000 m heights, since the CO₂ concentrations below 2,000 m could be highly influenced by airport pollution, and above 6,000 m CO₂ are fully mixed. At the heights of 2,000–6,000 m, every 500 m was divided into one layer, and in each layer, the observations were averaged and compared with the simulations.

Two forward simulations from May 2009 to December 2014 using the MOZART-4 model and the prior and posterior fluxes of the 12 TBMs were conducted to create prior and posterior CO₂ concentrations, respectively. The initial field at 00:00 UTC 01 May 2009 was obtained from the reanalysis concentration of Carbon Tracker CT2019B (CT2019B) (Jacobson et al., 2020). The mean deviation (BIAS) and root mean square error (RMSE) were used as reference indicators for the evaluation results. The monthly mean BIAS and RMSE at each continent were calculated.

3. Results

3.1. Uncertainty Reductions on Different Spatial Scales

As mentioned in Methods, 12 TBMs were used as prior fluxes. The NEE of these 12 TBMs has large differences. On the global scale, the mean annual NEEs from 2011 to 2014 are in the range of -2.66 (CASA model) to -9.97 (LPX-Bern model) PgC yr⁻¹ (Figure S2 in Supporting Information S1). We treat the standard deviation of the 12 TBMs' NEE as the 1- σ uncertainty and the mean of the 12 TBMs as the best estimate of NEE for one region. To explore the spatial scales at which GOSAT XCO₂ retrievals can provide robust NEE estimates, we analyze the relative prior uncertainty and uncertainty reduction rate (URR) after constraints at the global scale, the hemispheric scale (northern mid to high latitudes, tropical latitudes, southern middle latitudes), the continental scale, the half of continental scale (1/2 continent), the quarter of continental scale (1/4 continent), and small regions. The definition of the hemispheric scale and the latter three scales is given in Figure S1b–S1d in Supporting Information S1.

Figure 1 shows the relative uncertainties of the prior and posterior NEEs and their URRs after constraint using GOSAT XCO₂ on different spatial scales. Clearly, the relative prior uncertainty increases with decreasing spatial scale. On the global scale, the relative prior uncertainty is about 40%; on the continent, 1/2 continent, and 1/4 continent scales, the mean relative prior uncertainties are 47%, 53%, and 54%, respectively. On small regions (51 regions for global land, same thereafter), the mean relative prior uncertainty reaches 61%, with a range from 29% to 345%, and Figure S3 in Supporting Information S1 presents relative prior uncertainty views for small regions. The continent-scale relative prior uncertainty ranges from 36% to 88%.

After being constrained by the GOSAT XCO₂ retrievals, the uncertainty of the posterior NEE is substantially reduced. We find that the URR is significantly related to the spatial scale. The larger the spatial scale, the larger the URR, and vice versa. From the whole globe to the mean of 51 regions, the URR decreases from 95% to 16%. On the continental scale, the mean URR is 57%. N. America has the largest URR, with a value of 81%, followed by Asia (73%), S. America (58%), Africa (48%), and Australia (42%), and Europe has the smallest URR, with a value of only 39%. On small regions, posterior uncertainty decreased in most regions (0%–54%), except for 6 regions (located in northern Asia, eastern North America, Amazonia, and Southeast Asia) where posterior uncertainty increased to some extent (3%–43%), which may be related to the settings of prior uncertainty and observation errors (Figure S4 in Supporting Information S1). Moreover, the relative posterior uncertainty is lower than the prior on global to 1/4 continental scales, while in small regions, the relative posterior uncertainty outweighs the prior. This suggests that the GOSAT XCO₂ retrievals can constrain the terrestrial's NEE well at the continental scale, but has limited ability to constrain carbon fluxes at subcontinental or smaller scales, implying that the inversion results on sub-continental scales are highly related to the adopted prior NEE.

3.2. Annual and Seasonal Cycles of NEE on the Continental Scale

As mentioned above, on the continental scale, the posterior fluxes converge significantly. For prior fluxes, in Asia, N. America, Europe, S. America, Africa, and Australia, their averaged NEEs during the study period from the 12




Figure 1. Uncertainty at different scales and terrestrial carbon sink on the continental scale. (a) Relative uncertainties of the prior and posterior fluxes and uncertainty reduction rates after constrained using GOSAT XCO₂ in different spatial scales, and (b) annual prior and posterior net ecosystem exchanges (NEEs) on the global and continental scales. The uncertainty is depicted as the standard deviation of the simulated NEEs by the 12 terrestrial biosphere models.

TBMs are in the range of -0.25 to -3.27 , -0.13 to -1.76 , -0.24 to -1.26 , -0.29 to -1.62 , -0.31 to -2.14 , and -0.01 to -0.69 PgC yr⁻¹, with mean of -2.00 ± 0.91 , -0.99 ± 0.50 , -0.70 ± 0.34 , -1.10 ± 0.40 , -1.20 ± 0.58 , and -0.23 ± 0.20 PgC yr⁻¹, respectively. After constraining using XCO₂ retrievals, we obtain the mean NEEs of -2.07 ± 0.25 , -0.85 ± 0.09 , -0.76 ± 0.21 , -0.36 ± 0.17 , -0.36 ± 0.30 , and -0.15 ± 0.11 PgC yr⁻¹, respectively (Table S2 in Supporting Information S1).

We further explore whether the seasonal cycles of continental-scale NEE also converged significantly. As shown in Figure 2, for the prior fluxes, the monthly NEEs of different TBMs vary largely in all continents. In Asia, Europe, and N. America, although all models show strong land carbon sinks in warm seasons (May–September), and clear land carbon sources during the cold seasons, however, the seasonal magnitudes vary significantly across models, which are in the range of 0.39 – 2.88 PgC mo⁻¹, 0.29 – 1.41 PgC mo⁻¹, and 0.17 – 1.92 PgC mo⁻¹, respectively, with corresponding mean seasonal magnitudes of 1.34 ± 0.62 , 0.81 ± 0.26 , 0.96 ± 0.44 PgC mo⁻¹. Moreover, in Africa, S. America, and Australia, the different TBMs show very inconsistent seasonal cycles. For example, in Australia, some models show carbon sinks from April to October, some models show the opposite,




Figure 2. Averaged prior and posterior seasonal cycle of net ecosystem exchange (NEE) in different continents during 2011–2014. The lighter lines correspond to the NEEs of different terrestrial biosphere models, and the darker lines represent the multiple models mean.

and there are individual models that show carbon sinks throughout the year. The mean seasonal magnitudes of Africa, S. America, and Australia are 0.38 ± 0.13 , 0.51 ± 0.29 , and 0.19 ± 0.16 PgC mo⁻¹, respectively.

For the posterior fluxes, the seasonal cycles of different TBMs are in a narrow spread. Compared to the prior magnitudes, the posterior magnitudes have increased in Asia, N. America, Europe, and Africa, with Africa in particular more than doubling, while in S. America and Australia, they have decreased. The mean seasonal magnitudes of Asia, N. America, Europe, Africa, S. America, and Australia are 1.33 ± 0.27 , 1.10 ± 0.16 , 0.92 ± 0.16 , 0.31 ± 0.08 , 0.45 ± 0.11 , and 0.16 ± 0.11 PgC mo⁻¹, respectively. Uncertainties of their magnitudes are reduced by a range from 15% to 64%. In addition to more unified amplitudes, basically, all TBMs also present a consistent phase in their seasonal cycle. Particularly, in the prior NEEs, there are individual models whose results deviate significantly from others. For example, in Asia, North America, and Europe, one model shows neutral results in summer, and in S. America and Australia, there is a model showing abnormally high sources in June–October. After constraint by GOSAT observations, these anomalies of individual patterns disappear.

When comparing the multi-model mean prior and posterior seasonal cycles, in Asia, Europe, North America and Australia, the posterior seasonal cycle is consistent with the prior results, but the carbon sink is stronger in summer and the carbon source is stronger in autumn. In Africa, South America, and Australia, the posterior and prior seasonal cycles are quite different. In Africa, the prior NEEs show carbon sinks throughout the year, with the strongest carbon sinks in July–August and the weakest sinks in February and November; while the posterior NEE shows that there are significant carbon sources from October to December, and the strongest carbon sinks in August–September. In South America, the prior NEEs show a unimodal distribution, with the strongest sink and source in January and September, respectively; but the posterior results show that the month with strongest source appear in October.

3.3. Evaluation for the Inversion Results

We evaluated the inversion results using independent surface CO₂ observations over the globe. Figure 3 shows the continental averaged monthly mean observed CO₂ concentrations and the 12 TBMs averaged prior and posterior CO₂ concentrations. Compared to the prior CO₂ concentrations, except for Asia, the posterior concentrations are much closer to the observed values over all continents. The RMSE between the observations and simulations in Europe, N. America, S. America, Africa, and Australia decrease from a priori of 2.15, 2.37, 2.22, 1.77, and 1.58 ppm to a posteriori of 1.60, 1.50, 0.81, 0.53, and 0.61 ppm, respectively, with reduction rates of RMSE in the range of 26% ~ 70%. For the individual models (Figure S5 in Supporting Information S1), the BIAS and RMSE




Figure 3. Time series of modeled and observed monthly mean CO₂ concentrations for (a), Asia, (b), Europe, (c), N. America, (d), S. America, (e), Africa, and (f), Oceania. The embedded map in the upper left corner shows the location of the stations used in each continent.

of the posterior CO₂ are also lower than those of the prior CO₂ for almost all models and in all the continents. Generally, the prior CO₂ of the LPX-Bern, ORCHIDEE, ORCHIDEEv3, SDGVM, and VISIT models have larger RMSE and BIAS than the other models in all continents. After being constrained with XCO₂ data, the posterior CO₂ RMSE of these five models are similar with those of the others. In Asia, for the prior CO₂ concentrations, there are about half of the models with negative biases, and the rest with positive biases, with values in the range of $-3.19 \sim 5.56$ ppm, which results in a very small BIAS in the mean prior CO₂ of 0.17 ppm, while for the posterior CO₂, almost all models have small positive biases, with values in the range of $-0.07 \sim 2.68$ ppm and average bias of 1.23 ppm. In Southeast Asia, compared with the aircraft observations, the prior CO₂ have large negative bias (about -2.5 ppm), while the posterior CO₂ have a much smaller bias, with a value about -0.5 ppm (Figure S6 in Supporting Information S1). This indicates that the inversion results in Asia of all TBMs are also improved.

It can be found that the posterior CO₂ in Asia agrees well with the observation in summer, but in winter, the posterior concentration is higher than the observation, indicating that the carbon source in Asia was overestimated in winter. Although the posterior concentrations in N. America and Europe match the observations better overall, similar characteristics to Asia were observed, that is, the differences between the posterior concentrations and the observations are greater in winter than in summer, suggesting it might be caused by difficulties representing transport in the stable boundary layer in winter (Figure S7 in Supporting Information S1).

4. Discussion and Conclusion

With NEEs from 12 different TBMs, our work produces a robust estimate at the continental scale using GOSAT XCO₂, with very consistent annual mean carbon fluxes and seasonal cycles. The assessment of the results by

independent observations shows that the posterior concentrations are closer to the observations. Compared to previous estimates, the estimated net biosphere exchanges (NBE, =NEE + FIRE) in N. America, Europe, S. America, and Africa are close to or between the estimates of GCAS2021 (Jiang et al., 2022) and CMS-Flux NBE 2020 (Liu et al., 2021) during the same period (Figure S8 in Supporting Information S1), which were inferred from the same satellite retrievals as this study; while in Asia, the NBE of this study is significantly stronger than both, and in Australia, the sink in this study is weak. In N. America, and Europe our results are in the range of the NBEs constrained using surface measurements, namely CT2019B (Jacobson et al., 2020), Jena CarboScope s10oc_v2020 (Rodenbeck et al., 2018), and CAMS v18r2 (Chevallier et al., 2010) (Figure S8 in Supporting Information S1). In Asia our results are close to the estimate of Jena CarboScope s10oc_v2020 but significantly stronger than the other two estimates. In S. America we show a stronger sink, while in Africa, we show a stronger source compared to all three estimates. Compared to the other studies, Asia's NEE is also comparable to the estimate of Zhang et al. (2014), who used Asia's ground and aircraft observations as many as possible, and less than the estimate based on eddy covariance measurements (Ichii et al., 2017). Compared to the state-of-the-art bottom up estimate for the period of 2000–2009 (Ciais et al., 2021), this study shows a stronger sink in N. America, EuroAsia, and S. America, but a weaker one in Africa and Australia. On globe land, the NEE is reduced from a priori of -6.22 ± 2.48 PgC yr⁻¹ to a posteriori of -4.55 ± 0.12 PgC yr⁻¹. Combined with the prescribed fluxes of ocean (-2.45 PgC yr⁻¹), fire (1.93 PgC yr⁻¹), and fossil fuel and cement (9.68 PgC yr⁻¹), the posterior global net flux to the atmosphere is 4.61 PgC yr⁻¹, which is very close to the observed mean atmospheric CO₂ growth rate of 4.51 PgC yr⁻¹ (Friedlingstein et al., 2020).

The setting of prior uncertainties and observation errors, and the spatial resolution of observation aggregation can affect the estimates of NEE. When using the standard deviations of the 12 TBMs as the prior uncertainties in each region (Case Q), the URRs in most regions of high latitudes are lower than those of the Base case. The reason is that with this scheme, the given uncertainty for each prior flux at high latitude regions is greater (Figure S9 in Supporting Information S1). Besides, we also find a significant increase of URR in the tropical regions of Amazon and Indochina, indicating that these regions are sensitive to a priori uncertainty settings. The evaluation of inversions using aircraft observations in the Amazon by Wang et al. (2018) also suggests that reliable interannual fluxes may not be available in the Amazon at present. On the continental scale, the URRs decrease on all continents except Europe, while the estimated NEE in all continents does not change much compared with the Base case (Figure S10 in Supporting Information S1). When using a uniform and much smaller observation error (Case R, as described in “Method”), the URRs have increased in most regions, especially at high latitudes (Figure S4 in Supporting Information S1). When increasing the spatial resolution of observation aggregation, as expected, the URRs increased in most regions as well, especially when the resolution was increased to $5^\circ \times 5^\circ$ (Figure S4 in Supporting Information S1). The seasonal cycles of continental-scale NEEs are also more converged across TBMs (Figure S11 in Supporting Information S1). For the estimates of NEE, at the global scale, Case Q, Case R, Case A10, and Case A5 have almost the same annual carbon sinks with Base Case. At the continental scale, although the inverted NEEs of different schemes differ slightly, the differences between schemes are basically significantly smaller than the differences between the a priori and the a posteriori (Figure S13, Table S2 in Supporting Information S1). Overall, the different prior uncertainty and observation error settings and different spatial resolution of observation aggregation do not change the situation that URR decreases significantly as the spatial scale decreases, but with smaller observation error and higher aggregation resolution, the decline rates are reduced. With different schemes, the mean URRs below the continental scale are all less than 0.5 (Figure S12 in Supporting Information S1). We conclude that currently, the GOSAT XCO₂ can only give a robust estimate of the carbon flux on the continental scale, and as the research by Takagi et al. (2014), under the current satellite observing capacity, improving the XCO₂ accuracy can effectively reduce the spatial scale of robust carbon flux estimates.

Conflict of Interest

The authors declare no conflicts of interest relevant to this study.

Data Availability Statement

The V9 ACOS L2 data products of GOSAT XCO₂ column concentrations is publicly available at https://oco2.gesdisc.eosdis.nasa.gov/data/GOSAT_TANSO_Level2/ACOS_L2_Lite_FP.9r. The TRENDY TBMs data are available at <https://sites.exeter.ac.uk/trendy>. The CarbonTracker CT2017 fluxes used as prior information in

the model simulations can be accessed from the website <https://gml.noaa.gov/aftp/products/carbontracker/co2/CT2017>. The CarbonTracker CT2019B results provided by NOAA Earth System Research Laboratories (ESRL), Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov>. The ObsPack data can be downloaded from <https://gml.noaa.gov/ccgg/obspack/data.php>. The CONTRAIL data can be obtained from the ObsPack data set. The monthly carbon fluxes for the 51 terrestrial regions estimated in this study are available at <https://doi.org/10.5281/zenodo.7520065>.

Acknowledgments

This work is supported by the National Key R&D Program of China (Grant 2021YFB3901001), Fengyun Application Pioneering Project (Grant FY-APP-2022.0505), the Research Funds for the Frontiers Science Center for Critical Earth Material Cycling, Nanjing University (Grant 090414380031), and the National Natural Science Foundation of China (Grant 41907378). We acknowledge all atmospheric data providers to obspack_CO2_1_GLOBALVIEWplus_v7.0_2021-08-18. We acknowledge the CONTRAIL project team members for the public availability of their data as part of the ObsPack data package. The GOSAT data are produced by the ACOS/OCO-2 project at the Jet Propulsion Laboratory, California Institute of Technology, and obtained from the JPL website, CO2.jpl.nasa.gov. We acknowledge the members of the GOSAT Project for their contribution to this work. We are also grateful to the High-Performance Computing Center (HPCC) of Nanjing University for doing the numerical calculations in this paper on its blade cluster system.

References

- Baker, D. F., Bösch, H., Doney, S. C., O'Brien, D., & Schimel, D. S. (2010). Carbon source/sink information provided by column CO₂ measurements from the Orbiting Carbon Observatory. *Atmospheric Chemistry and Physics*, 10(9), 4145–4165. <https://doi.org/10.5194/acp-10-4145-2010>
- Baker, D. F., Law, R. M., Gurney, K. R., Rayner, P., Peylin, P., Denning, A. S., et al. (2006). TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO₂ fluxes, 1988–2003. *Global Biogeochemical Cycles*, 20(1). <https://doi.org/10.1029/2004GB002439>
- Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., et al. (2013). Global CO₂ fluxes estimated from GOSAT retrievals of total column CO₂. *Atmospheric Chemistry and Physics*, 13(17), 8695–8717. <https://doi.org/10.5194/acp-13-8695-2013>
- Buitenhuis, E., Le Quere, C., Aumont, O., Beaugrand, G., Bunker, A., Hirst, A., et al. (2006). Biogeochemical fluxes through mesozooplankton. *Global Biogeochemical Cycles*, 20(2). <https://doi.org/10.1029/2005GB002511>
- Byrne, B., Jones, D. B. A., Strong, K., Polavarapu, S. M., Harper, A. B., Baker, D. F., & Maksyutov, S. (2019). On what scales can GOSAT flux inversions constrain anomalies in terrestrial ecosystems? *Atmospheric Chemistry and Physics*, 19(20), 13017–13035. <https://doi.org/10.5194/acp-19-13017-2019>
- Byrne, B., Liu, J., Lee, M., Baker, I., Bowman, K. W., Deutscher, N. M., et al. (2020). Improved constraints on northern extratropical CO₂ fluxes obtained by combining surface-based and space-based atmospheric CO₂ measurements. *Journal of Geophysical Research-Atmospheres*, 125(15), e2019JD032029. <https://doi.org/10.1029/2019JD032029>
- Chen, J. M., Liu, J., Cihlar, J., & Goulden, M. L. (1999). Daily canopy photosynthesis model through temporal and spatial scaling for remote sensing applications. *Ecological Modelling*, 124(2–3), 99–119. [https://doi.org/10.1016/S0304-3800\(99\)00156-8](https://doi.org/10.1016/S0304-3800(99)00156-8)
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., et al. (2010). CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements. *Journal of Geophysical Research*, 115(D21), D21307. <https://doi.org/10.1029/2010JD013887>
- Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., & Bousquet, P. (2014). Toward robust and consistent regional CO₂ flux estimates from in situ and spaceborne measurements of atmospheric CO₂. *Geophysical Research Letters*, 41(3), 1065–1070. <https://doi.org/10.1002/2013GL058772>
- Chevallier, F., Remaud, M., O'Dell, C. W., Baker, D., Peylin, P., & Cozic, A. (2019). Objective evaluation of surface- and satellite-driven carbon dioxide atmospheric inversions. *Atmospheric Chemistry and Physics*, 19(22), 14233–14251. <https://doi.org/10.5194/acp-19-14233-2019>
- Ciais, P., Yao, Y., Gasser, T., Baccini, A., Wang, Y., Lauerwald, R., et al. (2021). Empirical estimates of regional carbon budgets imply reduced global soil heterotrophic respiration. *National Science Review*, 8(2). <https://doi.org/10.1093/nsr/nwaa145>
- Cox, A., Di Sarra, A. G., Vermeulen, A., Manning, A., Beyersdorf, A., Zahn, A., et al. (2021). Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957–2020; obspack_CO2_1_GLOBALVIEWplus_v7.0_2021-08-18. Retrieved from: http://www.esrl.noaa.gov/gmd/ccgg/obspack/data.php?id=obspack_CO2_1_GLOBALVIEWplus_v7.0_2021-08-18
- Crowell, S., Baker, D., Schuh, A., Basu, S., Jacobson, A. R., Chevallier, F., et al. (2019). The 2015–2016 carbon cycle as seen from OCO-2 and the global in situ network. *Atmospheric Chemistry and Physics*, 19(15), 9797–9831. <https://doi.org/10.5194/acp-19-9797-2019>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597. <https://doi.org/10.1002/qj.828>
- Deng, F., & Chen, J. M. (2011). Recent global CO₂ flux inferred from atmospheric CO₂ observations and its regional analyses. *Biogeosciences*, 8(11), 3263–3281. <https://doi.org/10.5194/bg-8-3263-2011>
- Deng, F., Jones, D. B. A., Henze, D. K., Bousseret, N., Bowman, K. W., Fisher, J. B., et al. (2014). Inferring regional sources and sinks of atmospheric CO₂ from GOSAT XCO₂ data. *Atmospheric Chemistry and Physics*, 14(7), 3703–3727. <https://doi.org/10.5194/acp-14-3703-2014>
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., et al. (2010). Description and evaluation of the model for Ozone and related chemical Tracers, version 4 (MOZART-4). *Geoscientific Model Development*, 3(1), 43–67. <https://doi.org/10.5194/gmd-3-43-2010>
- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., et al. (2020). Global carbon budget 2020. *Earth System Science Data*, 12(4), 3269–3340. <https://doi.org/10.5194/essd-12-3269-2020>
- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., et al. (2002). Towards robust regional estimates of CO₂ sources and sinks using atmospheric transport models. *Nature*, 415(6872), 626–630. <https://doi.org/10.1038/415626a>
- Haverd, V., Smith, B., Nieradzik, L., Briggs, P. R., Woodgate, W., Trudinger, C. M., et al. (2018). A new version of the CABLE land surface model (Subversion revision 14601) incorporating land use and land cover change, woody vegetation demography, and a novel optimisation-based approach to plant coordination of photosynthesis. *Geoscientific Model Development*, 11(7), 2995–3026. <https://doi.org/10.5194/gmd-11-2995-2018>
- Houweling, S., Baker, D., Basu, S., Boesch, H., Butz, A., Chevallier, F., et al. (2015). An intercomparison of inverse models for estimating sources and sinks of CO₂ using GOSAT measurements. *Journal of Geophysical Research: Atmospheres*, 120(10), 5253–5266. <https://doi.org/10.1002/2014jd022962>
- Houweling, S., Breon, F. M., Aben, I., Rodenbeck, C., Gloor, M., Heimann, M., & Ciais, P. (2004). Inverse modeling of CO₂ sources and sinks using satellite data: A synthetic inter-comparison of measurement techniques and their performance as a function of space and time. *Atmospheric Chemistry and Physics*, 4(2), 523–538. <https://doi.org/10.5194/acp-4-523-2004>
- Ichii, K., Ueyama, M., Kondo, M., Saigusa, N., Kim, J., Alberto, M. C., et al. (2017). New data-driven estimation of terrestrial CO₂ fluxes in Asia using a standardized database of eddy covariance measurements, remote sensing data, and support vector regression. *Journal of Geophysical Research: Biogeosciences*, 122(4), 767–795. <https://doi.org/10.1002/2016JG003640>
- Jacobson, A. R., Schuldt, K. N., Miller, J. B., Oda, T., Tans, P., Arlyn, A., et al. (2020). *CarbonTracker CT2019B*. NOAA Global Monitoring Laboratory.

- Jiang, F., Ju, W., He, W., Wu, M., Wang, H., Wang, J., et al. (2022). A 10-year global monthly averaged terrestrial net ecosystem exchange dataset inferred from the ACOS GOSAT v9 XCO₂ retrievals (GCAS2021). *Earth System Science Data*, 14(7), 3013–3037. <https://doi.org/10.5194/essd-14-3013-2022>
- Jiang, F., Wang, H., Chen, J. M., Ju, W., Tian, X., Feng, S., et al. (2021). Regional CO₂ fluxes from 2010 to 2015 inferred from GOSAT XCO₂ retrievals using a new version of the Global Carbon Assimilation System. *Atmospheric Chemistry and Physics*, 21(3), 1963–1985. <https://doi.org/10.5194/acp-21-1963-2021>
- Jiang, F., Wang, H. W., Chen, J. M., Zhou, L. X., Ju, W. M., Ding, A. J., et al. (2013). Nested atmospheric inversion for the terrestrial carbon sources and sinks in China. *Biogeosciences*, 10(8), 5311–5324. <https://doi.org/10.5194/bg-10-5311-2013>
- Ju, W. M., Chen, J. M., Black, T. A., Barr, A. G., Liu, J., & Chen, B. Z. (2006). Modelling multi-year coupled carbon and water fluxes in a boreal aspen forest. *Agricultural and Forest Meteorology*, 140(1–4), 136–151. <https://doi.org/10.1016/j.agrformet.2006.08.008>
- Kato, E., Kinoshita, T., Ito, A., Kawamiya, M., & Yamagata, Y. (2013). Evaluation of spatially explicit emission scenario of land-use change and biomass burning using a process-based biogeochemical model. *Journal of Land Use Science*, 8(1), 104–122. <https://doi.org/10.1080/1747423X.2011.628705>
- Lienert, S., & Joos, F. (2018). A Bayesian ensemble data assimilation to constrain model parameters and land-use carbon emissions. *Biogeosciences*, 15(9), 2909–2930. <https://doi.org/10.5194/bg-15-2909-2018>
- Liu, J., Baskaran, L., Bowman, K., Schimel, D., Bloom, A. A., Parazoo, N. C., et al. (2021). Carbon monitoring system flux net biosphere exchange 2020 (CMS-Flux NBE 2020). *Earth System Science Data*, 13(2), 299–330. <https://doi.org/10.5194/essd-13-299-2021>
- Lurton, T., Balkanski, Y., Bastrikov, V., Bekki, S., Bopp, L., Braconnot, P., et al. (2020). Implementation of the CMIP6 forcing data in the IPSL-CM6A-LR model. *Journal of Advances in Modeling Earth Systems*, 12(4), e2019MS001940. <https://doi.org/10.1029/2019MS001940>
- Machida, T., Matsueda, H., Sawa, Y., Nakagawa, Y., Hirotsu, K., Kondo, N., et al. (2008). Worldwide measurements of atmospheric CO₂ and other Trace gas species using commercial airlines. *Journal of Atmospheric and Oceanic Technology*, 25(10), 1744–1754. <https://doi.org/10.1175/2008JTECHA1082.1>
- Machida, T., Matsueda, H., Sawa, Y., Niwa, Y., Tsuboi, K., Katsumata, K., et al. (2019). *Atmospheric trace gas data from the CONTRAIL flask air sampling over the Eurasian continent, ver.2021.1.0*. Center for Global Environmental Research, NIES. <https://doi.org/10.17595/20190828.002> (Reference date: 2021/10/29)
- Maksyutov, S., Takagi, H., Valsala, V. K., Saito, M., Oda, T., Saeki, T., et al. (2013). Regional CO₂ flux estimates for 2009–2010 based on GOSAT and ground-based CO₂ observations. *Atmospheric Chemistry and Physics*, 13(18), 9351–9373. <https://doi.org/10.5194/acp-13-9351-2013>
- Matsueda, H., Machida, T., Sawa, Y., Nakagawa, Y., Hirotsu, K., Ikeda, H., et al. (2008). Evaluation of atmospheric CO₂ measurements from new flask air sampling of JAL airliner observations. *Papers in Meteorology and Geophysics*, 59, 1–17. <https://doi.org/10.2467/mripapers.59.1>
- Matsueda, H., Machida, T., Sawa, Y., & Niwa, Y. (2015). Long-term change of CO₂ latitudinal distribution in the upper troposphere. *Geophysical Research Letters*, 42(7), 2508–2514. <https://doi.org/10.1002/2014GL062768>
- Meiyappan, P., Jain, A. K., & House, J. I. (2015). Increased influence of nitrogen limitation on CO₂ emissions from future land use and land use change. *Global Biogeochemical Cycles*, 29(9), 1524–1548. <https://doi.org/10.1002/2015GB005086>
- Monteil, G., Broquet, G., Scholze, M., Lang, M., Karstens, U., Gerbig, C., et al. (2020). The regional European atmospheric transport inversion comparison, EUROCOM: First results on European-wide terrestrial carbon fluxes for the period 2006–2015. *Atmospheric Chemistry and Physics*, 20(20), 12063–12091. <https://doi.org/10.5194/acp-20-12063-2020>
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., et al. (2018). Improved retrievals of carbon dioxide from Orbiting Carbon Observatory-2 with the version 8 ACOS algorithm. *Atmospheric Measurement Techniques*, 11(12), 6539–6576. <https://doi.org/10.5194/amt-11-6539-2018>
- Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., et al. (2013). Global atmospheric carbon budget: Results from an ensemble of atmospheric CO₂ inversions. *Biogeosciences*, 10(10), 6699–6720. <https://doi.org/10.5194/bg-10-6699-2013>
- Philip, S., Johnson, M. S., Potter, C., Genovesse, V., Baker, D. F., Haynes, K. D., et al. (2019). Prior biosphere model impact on global terrestrial CO₂ fluxes estimated from OCO-2 retrievals. *Atmospheric Chemistry and Physics*, 19(20), 13267–13287. <https://doi.org/10.5194/acp-19-13267-2019>
- Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M., Mooney, H. A., & Klooster, S. A. (1993). Terrestrial ecosystem production - A process model-based on global satellite and surface data. *Global Biogeochemical Cycles*, 7(4), 811–841. <https://doi.org/10.1029/93GB02725>
- Rayner, P. J., Enting, I. G., Francey, R. J., & Langenfelds, R. (1999). Reconstructing the recent carbon cycle from atmospheric CO₂, δ¹³C and O₂/N₂ observations. *Tellus Series B Chemical and Physical Meteorology*, 51(2), 213–232. <https://doi.org/10.3402/tellusb.v51i2.16273>
- Rodenbeck, C., Houweling, S., Gloor, M., & Heimann, M. (2003). CO₂ flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport. *Atmospheric Chemistry and Physics*, 3(6), 1919–1964. <https://doi.org/10.5194/acp-3-1919-2003>
- Rodenbeck, C., Zaehle, S., Keeling, R., & Heimann, M. (2018). How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO₂ data. *Biogeosciences*, 15(8), 2481–2498. <https://doi.org/10.5194/bg-15-2481-2018>
- Sierk, B., Fernandez, V., Bézy, J. L., Meijer, Y., Durand, Y., Bazalgette Courrèges-Lacoste, G., et al. (2021). The Copernicus CO₂M mission for monitoring anthropogenic carbon dioxide emissions from space. In *International Conference on Space Optics—ICSO 2020*. Paper presented at the. <https://doi.org/10.1117/12.2599613>
- Takagi, H., Houweling, S., Andres, R. J., Belikov, D., Bril, A., Boesch, H., et al. (2014). Influence of differences in current GOSAT XCO₂ retrievals on surface flux estimation. *Geophysical Research Letters*, 41(7), 2598–2605. <https://doi.org/10.1002/2013gl059174>
- Taylor, T. E., O'Dell, C. W., Crisp, D., Kuze, A., Lindqvist, H., Wennberg, P. O., et al. (2022). An 11-year record of XCO₂ estimates derived from GOSAT measurements using the NASA ACOS version 9 retrieval algorithm. *Earth System Science Data*, 14(1), 325–360. <https://doi.org/10.5194/essd-14-325-2022>
- Thompson, R. L., Patra, P. K., Chevallier, F., Maksyutov, S., Law, R. M., Ziehn, T., et al. (2016). Top-down assessment of the Asian carbon budget since the mid 1990s. *Nature Communications*, 7(1), 10724. <https://doi.org/10.1038/ncomms10724>
- Tian, H., Chen, G., Lu, C., Xu, X., Hayes, D. J., Ren, W., et al. (2015). North American terrestrial CO₂ uptake largely offset by CH₄ and N₂O emissions: Toward a full accounting of the greenhouse gas budget. *Climatic Change*, 129(3–4), 413–426. <https://doi.org/10.1007/s10584-014-1072-9>
- Vuichard, N., Messina, P., Luyssaert, S., Guenet, B., Zaehle, S., Ghattas, J., et al. (2019). Accounting for carbon and nitrogen interactions in the global terrestrial ecosystem model ORCHIDEE (trunk version, rev 4999): Multi-scale evaluation of gross primary production. *Geoscientific Model Development*, 12(11), 4751–4779. <https://doi.org/10.5194/gmd-12-4751-2019>
- Walker, A. P., Quaife, T., van Bodegom, P. M., De Kauwe, M. G., Keenan, T. F., Joiner, J., et al. (2017). The impact of alternative trait-scaling hypotheses for the maximum photosynthetic carboxylation rate (V_{cmax}) on global gross primary production. *New Phytologist*, 215(4), 1370–1386. <https://doi.org/10.1111/nph.14623>

- Wang, H., Jiang, F., Liu, Y., Yang, D., Wu, M., He, W., et al. (2022). Global terrestrial ecosystem carbon flux inferred from TanSat XCO₂ retrievals. *Journal of Remote Sensing*, 2022, 1–12. <https://doi.org/10.34133/2022/9816536>
- Wang, H., Jiang, F., Wang, J., Ju, W., & Chen, J. M. (2019). Terrestrial ecosystem carbon flux estimated using GOSAT and OCO-2 XCO₂ retrievals. *Atmospheric Chemistry and Physics*, 19(18), 12067–12082. <https://doi.org/10.5194/acp-19-12067-2019>
- Wang, J., Jiang, F., Wang, H., Qiu, B., Wu, M., He, W., et al. (2021). Constraining global terrestrial gross primary productivity in a global carbon assimilation system with OCO-2 chlorophyll fluorescence data. *Agricultural and Forest Meteorology*, 304–305, 304–305. <https://doi.org/10.1016/j.agrformet.2021.108424>
- Wang, J. S., Kawa, S. R., Collatz, G. J., Sasakawa, M., Gatti, L. V., Machida, T., et al. (2018). A global synthesis inversion analysis of recent variability in CO₂ fluxes using GOSAT and in situ observations. *Atmospheric Chemistry and Physics*, 18(15), 11097–11124. <https://doi.org/10.5194/acp-18-11097-2018>
- Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., et al. (2017). Comparisons of the orbiting carbon observatory-2 (OCO-2) XCO₂ measurements with TCCON. *Atmospheric Measurement Techniques*, 10(6), 2209–2238. <https://doi.org/10.5194/amt-10-2209-2017>
- Yue, X., & Unger, N. (2015). The yale interactive terrestrial biosphere model version 1.0: Description, evaluation and implementation into NASA GISS ModelE2. *Geoscientific Model Development*, 8(8), 2399–2417. <https://doi.org/10.5194/gmd-8-2399-2015>
- Zaehle, S., & Friend, A. D. (2010). Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates. *Global Biogeochemical Cycles*, 24. <https://doi.org/10.1029/2009gb003521>
- Zhang, H. F., Chen, B. Z., van der Laan-Luijk, I. T., Machida, T., Matsueda, H., Sawa, Y., et al. (2014). Estimating Asian terrestrial carbon fluxes from CONTRAIL aircraft and surface CO₂ observations for the period 2006–2010. *Atmospheric Chemistry and Physics*, 14(11), 5807–5824. <https://doi.org/10.5194/acp-14-5807-2014>

References From the Supporting Information

- Connor, B. J., Boesch, H., Toon, G., Sen, B., Miller, C., & Crisp, D. (2008). Orbiting carbon observatory: Inverse method and prospective error analysis. *Journal of Geophysical Research*, 113(D5). <https://doi.org/10.1029/2006JD008336>